



early-onset glaucoma

Glaucoma is a group of eye disorders in which the optic nerves connecting the eyes and the brain are progressively damaged. This damage can lead to reduction in side (peripheral) vision and eventual blindness. Other signs and symptoms may include bulging eyes, excessive tearing, and abnormal sensitivity to light (photophobia). The term "early-onset glaucoma" may be used when the disorder appears before the age of 40.

In most people with glaucoma, the damage to the optic nerves is caused by increased pressure within the eyes (intraocular pressure). Intraocular pressure depends on a balance between fluid entering and leaving the eyes.

Usually glaucoma develops in older adults, in whom the risk of developing the disorder may be affected by a variety of medical conditions including high blood pressure (hypertension) and diabetes mellitus, as well as family history. The risk of early-onset glaucoma depends mainly on heredity.

Structural abnormalities that impede fluid drainage in the eye may be present at birth and usually become apparent during the first year of life. Such abnormalities may be part of a genetic disorder that affects many body systems, called a syndrome. If glaucoma appears before the age of 5 without other associated abnormalities, it is called primary congenital glaucoma.

Other individuals experience early onset of primary open-angle glaucoma, the most common adult form of glaucoma. If primary open-angle glaucoma develops during childhood or early adulthood, it is called juvenile open-angle glaucoma.

Frequency

Primary congenital glaucoma affects approximately 1 in 10,000 people. Its frequency is higher in the Middle East. Juvenile open-angle glaucoma affects about 1 in 50,000 people. Primary open-angle glaucoma is much more common after the age of 40, affecting about 1 percent of the population worldwide.

Genetic Changes

Approximately 10 percent to 33 percent of people with juvenile open-angle glaucoma have mutations in the *MYOC* gene. *MYOC* gene mutations have also been detected in some people with primary congenital glaucoma. The *MYOC* gene provides instructions for producing a protein called myocilin. Myocilin is found in certain structures of the eye, called the trabecular meshwork and the ciliary body, that regulate the intraocular pressure.

Researchers believe that myocilin functions together with other proteins as part of a protein complex. Mutations may alter the protein in such a way that the complex cannot be formed. Defective myocilin that is not incorporated into functional complexes may accumulate in the trabecular meshwork and ciliary body. The excess protein may prevent sufficient flow of fluid from the eye, resulting in increased intraocular pressure and causing the signs and symptoms of early-onset glaucoma.

Between 20 percent and 40 percent of people with primary congenital glaucoma have mutations in the *CYP1B1* gene. *CYP1B1* gene mutations have also been detected in some people with juvenile open-angle glaucoma. The *CYP1B1* gene provides instructions for producing a form of the cytochrome P450 protein. Like myocilin, this protein is found in the trabecular meshwork, ciliary body, and other structures of the eye.

It is not well understood how defects in the CYP1B1 protein cause signs and symptoms of glaucoma. Recent studies suggest that the defects may interfere with the early development of the trabecular meshwork. In the clear covering of the eye (the cornea), the CYP1B1 protein may also be involved in a process that regulates the secretion of fluid inside the eye. If this fluid is produced in excess, the high intraocular pressure characteristic of glaucoma may develop.

The CYP1B1 protein may interact with myocilin. Individuals with mutations in both the *MYOC* and *CYP1B1* genes may develop glaucoma at an earlier age and have more severe symptoms than do those with mutations in only one of the genes. Mutations in other genes may also be involved in early-onset glaucoma.

Inheritance Pattern

Early-onset glaucoma can have different inheritance patterns. Primary congenital glaucoma is usually inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. Most often, the parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but do not show signs and symptoms of the condition.

Juvenile open-angle glaucoma is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder. In some families, primary congenital glaucoma may also be inherited in an autosomal dominant pattern.

Other Names for This Condition

- hereditary glaucoma

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Glaucoma, congenital
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0020302/>
- Genetic Testing Registry: Primary open angle glaucoma juvenile onset 1
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1842028/>

Other Diagnosis and Management Resources

- GeneReview: Primary Congenital Glaucoma
<https://www.ncbi.nlm.nih.gov/books/NBK1135>
- MedlinePlus Encyclopedia: Glaucoma
<https://medlineplus.gov/ency/article/001620.htm>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Glaucoma
<https://medlineplus.gov/ency/article/001620.htm>
- Health Topic: Glaucoma
<https://medlineplus.gov/glaucoma.html>
- Health Topic: Optic Nerve Disorders
<https://medlineplus.gov/opticnervedisorders.html>

Additional NIH Resources

- National Eye Institute
https://nei.nih.gov/health/glaucoma/glaucoma_facts

Educational Resources

- American Optometric Association
<http://www.aoa.org/patients-and-public/eye-and-vision-problems/glossary-of-eye-and-vision-conditions/glaucoma>
- MalaCards: early-onset glaucoma
http://www.malacards.org/card/early_onset_glaucoma
- Merck Manual
<http://www.merckmanuals.com/home/eye-disorders/glaucoma/glaucoma>
- Orphanet: Hereditary glaucoma
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=359

Patient Support and Advocacy Resources

- Children's Glaucoma Foundation
http://www.childrensglaucomafoundation.org/_structure.php
- EyeSmart: What is Glaucoma?
<https://www.aaio.org/eye-health/diseases/what-is-glaucoma>
- Glaucoma Foundation
<http://www.glaucomafoundation.org/index.php>
- Glaucoma Research Foundation
<http://www.glaucoma.org/glaucoma/childhood-glaucoma-1.php>
- Glaucoma Service Foundation to Prevent Blindness
<http://willsglaucoma.org/>
- International Glaucoma Association
<http://www.glaucoma-association.com/>
- Pediatric Glaucoma and Cataract Family Association
<http://pgcfa.org/>

GeneReviews

- Primary Congenital Glaucoma
<https://www.ncbi.nlm.nih.gov/books/NBK1135>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22glaucoma%22+OR+%22early-onset+glaucoma%22>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28early-onset+glaucoma%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- GLAUCOMA 1, OPEN ANGLE, A
<http://omim.org/entry/137750>
- GLAUCOMA 3, PRIMARY CONGENITAL, A
<http://omim.org/entry/231300>

Sources for This Summary

- Achary MS, Reddy AB, Chakrabarti S, Panicker SG, Mandal AK, Ahmed N, Balasubramanian D, Hasnain SE, Nagarajaram HA. Disease-causing mutations in proteins: structural analysis of the CYP1B1 mutations causing primary congenital glaucoma in humans. *Biophys J*. 2006 Dec 15; 91(12):4329-39. Epub 2006 Sep 8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16963504>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1779944/>
- Acharya M, Mookherjee S, Bhattacharjee A, Bandyopadhyay AK, Daulat Thakur SK, Bhaduri G, Sen A, Ray K. Primary role of CYP1B1 in Indian juvenile-onset POAG patients. *Mol Vis*. 2006 Apr 20;12:399-404.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16688110>
- Bayat B, Yazdani S, Alavi A, Chiani M, Chitsazian F, Tusi BK, Suri F, Narooie-Nejhad M, Sanati MH, Elahi E. Contributions of MYOC and CYP1B1 mutations to JOAG. *Mol Vis*. 2008 Mar 13;14: 508-17.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18385784>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2268862/>
- Chakrabarti S, Kaur K, Rao KN, Mandal AK, Kaur I, Parikh RS, Thomas R. The transcription factor gene FOXC1 exhibits a limited role in primary congenital glaucoma. *Invest Ophthalmol Vis Sci*. 2009 Jan;50(1):75-83. doi: 10.1167/iops.08-2253. Epub 2008 Aug 15.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18708620>
- Chavarria-Soley G, Sticht H, Aklillu E, Ingelman-Sundberg M, Pasutto F, Reis A, Rautenstrauss B. Mutations in CYP1B1 cause primary congenital glaucoma by reduction of either activity or abundance of the enzyme. *Hum Mutat*. 2008 Sep;29(9):1147-53. doi: 10.1002/humu.20786.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18470941>
- Chen Y, Jiang D, Yu L, Katz B, Zhang K, Wan B, Sun X. CYP1B1 and MYOC mutations in 116 Chinese patients with primary congenital glaucoma. *Arch Ophthalmol*. 2008 Oct;126(10):1443-7. doi: 10.1001/archophth.126.10.1443.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18852424>

- Choudhary D, Jansson I, Sarfarazi M, Schenkman JB. Characterization of the biochemical and structural phenotypes of four CYP1B1 mutations observed in individuals with primary congenital glaucoma. *Pharmacogenet Genomics*. 2008 Aug;18(8):665-76. doi: 10.1097/FPC.0b013e3282ff5a36.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18622259>
- Firasat S, Riazuddin SA, Khan SN, Riazuddin S. Novel CYP1B1 mutations in consanguineous Pakistani families with primary congenital glaucoma. *Mol Vis*. 2008;14:2002-9. Epub 2008 Nov 3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18989382>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2579935/>
- OMIM: GLAUCOMA 1, OPEN ANGLE, A
<http://omim.org/entry/137750>
- OMIM: GLAUCOMA 3, PRIMARY CONGENITAL, A
<http://omim.org/entry/231300>
- GeneReview: Primary Congenital Glaucoma
<https://www.ncbi.nlm.nih.gov/books/NBK1135>
- Joe MK, Sohn S, Hur W, Moon Y, Choi YR, Kee C. Accumulation of mutant myocilins in ER leads to ER stress and potential cytotoxicity in human trabecular meshwork cells. *Biochem Biophys Res Commun*. 2003 Dec 19;312(3):592-600.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14680806>
- Kaur K, Reddy AB, Mukhopadhyay A, Mandal AK, Hasnain SE, Ray K, Thomas R, Balasubramanian D, Chakrabarti S. Myocilin gene implicated in primary congenital glaucoma. *Clin Genet*. 2005 Apr;67(4):335-40.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15733270>
- Ray K, Mukhopadhyay A, Acharya M. Recent advances in molecular genetics of glaucoma. *Mol Cell Biochem*. 2003 Nov;253(1-2):223-31. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14619973>
- Turalba AV, Chen TC. Clinical and genetic characteristics of primary juvenile-onset open-angle glaucoma (JOAG). *Semin Ophthalmol*. 2008 Jan-Feb;23(1):19-25. doi: 10.1080/08820530701745199. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18214788>
- Vasiliou V, Gonzalez FJ. Role of CYP1B1 in glaucoma. *Annu Rev Pharmacol Toxicol*. 2008;48:333-58. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17914928>
- Vincent AL, Billingsley G, Buys Y, Levin AV, Priston M, Trope G, Williams-Lyn D, Héon E. Digenic inheritance of early-onset glaucoma: CYP1B1, a potential modifier gene. *Am J Hum Genet*. 2002 Feb;70(2):448-60. Epub 2002 Jan 3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11774072>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC384919/>
- Weisschuh N, Schiefer U. Progress in the genetics of glaucoma. *Dev Ophthalmol*. 2003;37:83-93. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12876831>

- Wiggs JL, Lynch S, Ynagi G, Maselli M, Auguste J, Del Bono EA, Olson LM, Haines JL. A genomewide scan identifies novel early-onset primary open-angle glaucoma loci on 9q22 and 20p12. *Am J Hum Genet.* 2004 Jun;74(6):1314-20. Epub 2004 Apr 23.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15108121>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1182098/>
- Willoughby CE, Chan LL, Herd S, Billingsley G, Noordeh N, Levin AV, Buys Y, Trope G, Sarfarazi M, Héon E. Defining the pathogenicity of optineurin in juvenile open-angle glaucoma. *Invest Ophthalmol Vis Sci.* 2004 Sep;45(9):3122-30.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15326130>
- WuDunn D. Genetic basis of glaucoma. *Curr Opin Ophthalmol.* 2002 Apr;13(2):55-60. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11880716>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/condition/early-onset-glaucoma>

Reviewed: February 2009
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services